sdsJ qmilO 🛶

Deficiency in Infants, Children, and Adolescents.
Pediatrics, November 2008, 122(5)1142-1152.

Stegler E.E., Nelson S.E., Jeter J.M., Vitamin D supplementation of breastfed infants: a randomized dose-response trial Pediat Res 2014 Aug; 76(2): 177-183.

Mutritional Mickets. Horm Res Paediatr 2016;85:83-106. $^{\rm NW}$ and Vitamin D $^{\rm NW}$ Recommendations on Prevention and Management of (Flopai 'Te 10

*NRV – nutrient reference value.

10 hg (400 IU) 200%

Vitamin D (D3)

j capsale Nutrition information $NB\Lambda_*$

spell (gelatine, humectant - glycerol, water). [oil], cholecalciferol - vitamin D, capsule Ingredients: medium-chain triglycerides oil

physician before using this product. swallowed by the baby. The product should not be used in case of hypersensitivity to any of its ingredients. If you feed your baby vitamin D-fortified infant formula, consult your Marning: make sure that the capsule shell is not . The diet with vitamin D.

Recommended use: newborns and infants (up to 12 months): content of 1 twist-off capsule a day (400 IU of vitamin D3) or otherwise after consultation with physician. Administer with a meal. Twist and pull off

the tip of the capsule. In case of difficulties with opening the capsule, cut off the tip. The squeezed

content can be administered either directly into the

and partially breastled intants, to supplement infants (up to 12 months), especially for breastfed infants from birth to 12 months of age, 123

sunlight and age.
Vitamin D deficiency is common worldwide.
According to the opinion of experts, 400 IUV day (10 µg) or vitamin D is recommended for all instants from birth to 12 months of ene 123

sunscreen, clothing, amount of skin exposed to on a number of factors, including season of the year, skin pigmentation, latitude, use of of endogenous synthesis may vary depending The quantities of vitamin D produced as a resultbhosphorus.

normal blood calcium levels and promotes normal absorption/ utilisation of calcium and maintenance of normal teeth, helps to maintain function of the immune system in children. Furthermore, vitamin D contributes to the

needed for normal growth and development of bone in children and contributes to the normal Food supplement in the form of twist-off capsules containing vitamin D (D3). Vitamin D is observed for request based desirables and desirables are desirables.

yand editiv-blof

baby's mouth, poured in a spoon, or mixed with a spoonful of cold or room temperature soft food or expressed milk. Dispose the capsule shell. Do not exceed the recommended daily dose. supplements should not be used as a substitute for a varied and balanced diet. A varied and balanced diet and a healthy lifestyle are recommended.

30 twist-off capsules

Food supplement

For newborns and infants







Gold-Vit D₃ Baby Food supplement

Best before: see left side of the packaging.

12,8 g

Olimp

Olimp Laboratories Sp. z o.o. Pustynia 848, 92-200 Debica, Poland rel. +48 1468032200, fax +48 146803256 e-mail: info@olimpHabs.com, www.olimp-labs.com















425 mg